Johnson Johnson FAMILY OF COMPANIES

Project Name / Design GB_Benadryl Allergy One A Day_10mg Tablets_7s 14s 30s_PIL SNAS 3825

				51	,	· -	_		
Date	10.05.23					Country / Mark	et(s) - Langua	iges	English
Job / Ver	sion Numl	ber 40328	7857_r1			Technical Spec.	/ Drawing #	N/A	
Compon	ent Type	Leafle	t			Dimensions			
New Con	nponent /	ltem Code	XXXXXX			Manufacturing	/ Packaging F	ill Site	EN-UCB Farchim SA_Chemin de Cruix Blanche_Bulle_Switzerl
Old Com	ponent / h	tem Code	CIA79546	:		Printer Name	9061245		
Pharma	Code / Lae	tus Code	FP0			Print Method	Litho		
Product	/ Eurocode	e / Mat. ID C	ode 8714	206 8714406	73806	Substrate	white		
ArtWork	s ID: EM	IEA_2023_0	0092239_0	000		PC spec	PC-006331		

ARTWORK SPECIFICS	
PANTONE 287 C LITHOOFFSET	
REPRESENTATION: Colours represented with a diagonal line have b	een modified to aid PDF approval.
product. Schawk shall not be liable for any other damages and losses including but not limited to in delivery resulting from arrors or defects in any of Schawk's products. Client is solely responsibl language errors in the digital file, proofs, plates, files or other media to properly advise the end c	les sure that corrections are made before using this product. SchwkS liability is limited to fixing its own the costs of printing plates or cylinders, loss of print true, labo, schwsters, paints, other materials and delays e for (1) and failure on the file to comply with any applicable local, state or federal law or regulation, (10) and someme of any danger in the use or application of any modult can (1) and part of the life that violates the patent or otherwise. Printer is responsible for apply and ensure baccode and datamatrix code readability
	SGK is a Matthews International Corporation
FONTS USED	
HelveticaNeue MetaPlusBold-Roman	SMALLEST TEXT SIZE
MyriadPro-Regular	5.50 pt
Optima Optima-Bold	AVE. TEXT SIZE
Optima-BoldItalic	7.5 pt
ZapfDingbatsITC	





ONEADAY 10 mg TABLETS Cetirizine dihydrochloride

- This medicine is used to relieve the symptoms of hav fever and other allergic conditions such as
- pet or dust allergies and skin reactions. This medicine is for use by adults and children
- aged 12 years and over. A doctor must be contacted if symptoms worsen
- or do not improve after 3 days

Read the leaflet carefully before use. Keep this leaflet, you might need it again.

1 What the medicine is for

Benadryl Allergy One a Day 10 mg Tablets is a medicine which is used to relieve the symptoms of hav fever and other allergic conditions such as pet or dust allergies. The tablets contain 10 mg cetirizine dihydrochloride which is an antihistamine that helps relieve allergy symptoms such as sneezing, runny nose and watery eyes. It can also be used to treat allergic skin reactions. These include urticaria, also known as hives, where

the skin looks blotchy, with white raised wheals (bumps) surrounded by redness.

This medicine is for use in adults and children aged 12 years and over.

2 Before taking this medicine

This medicine is suitable for most adults and children aged 12 years and over, but a few people should not use it. If you are in any doubt, talk to your doctor or pharmacist.

X Do not take this medicine...

- If you have ever had a **bad reaction** to cetirizine dihydrochloride, to hydroxyzine or piperazine derivatives or any other ingredients of this medicine (See section 6). If you have serious kidney problems.
- **A** Talk to your doctor or pharmacist ■ If you are an epileptic or at risk of
- convulsions (fits)
- If this applies to you, **talk to a doctor or** pharmacist.

If you experience symptoms such as a sudden onset of fever, reddening of the skin with blisters containing pus, stop giving this medicine immediately and talk to a doctor or pharmacist (See section 4).

Warnings and precautions

If you are a patient with kidney problems, please ask your doctor for advice; if necessary, you will take a lower dose. The new dose will be determined by your doctor. If you have difficulty passing urine, you should ask your doctor for advice.

A If you are pregnant or breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

You should not take Benadryl Allergy One a Day 10 mg Tablets during breast feeding because cetirizine passes into breast milk.

0£

A Special warnings about this medicine

- In most people this product does not cause drowsiness. However, there are rare exceptions and you should take care when you use this product for the first time. If this product makes you feel drowsy, do not drive or operate machinery. This may be worsened by alcohol, sedatives or tranquilisers (used to treat anxiety or sleep disorders).
- As with all antihistamines, it is advisable to avoid alcohol consumption when taking this medicine.
- You should stop taking this product for 3 days before undergoing a skin allergy test because antihistamines can affect the results of the test. This product contains lactose. If you have been told by your doctor that you have an

intolerance to some sugars, contact your doctor before taking this medicinal product.

- **3** How to take this medicine For oral use only.
- Swallow with a glass of water.
- Do not take more than the stated dose shown in the table.

Children under 12 years old

This medicine is not recommended for children aged under 12 years old.

Adults and children aged 12 years and above

Age	Dose				
Adults and children aged 12 years and above	Take 1 tablet once a day				
 Do not take more than 1 tablet in 24 hours. If symptoms persist or worsen, stop use and consult your doctor or pharmacist 					

A If anyone has too much

If anyone has too much contact a doctor or your nearest Accident and Emergency department (Casualty) taking this leaflet and pack with you.

A If you forget to take the medicine

If you forget to take a dose, take the next dose when needed provided that you only take a maximum of 1 tablet in 24 hours. Do not take a double dose.

4 Possible side-effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

If you experience any of the following, stop taking this medicine and seek immediate medical help:

- Sudden onset of fever, reddening of the skin, or many small pustules (possible symptoms of Acute Generalised Exanthematous Pustulosis - AGEP) may occur within the first 2 days of treatment with this medicine (See section 2).
- Sudden signs of allergy such as rash, itching or hives on the skin, swelling of the face, lips, tongue or other parts of the body, shortness of breath, wheezing or trouble breathing
- Bruising or bleeding more easily than normal
- Convulsions (fits) or hallucinations
- Loss of consciousness

turn over 🕨

Johnson Johnson FAMILY OF COMPANIES

Project Name / Design GB_Benadryl Allergy One A Day_10mg Tablets_7s 14s 30s_PIL SNAS 3825

Date	10.05.23			Country / Market(s) - Languages			English
Job / Version Number 403287857_r1			Technical Spec	. / Drawing #	N/A		
Component Type Leaflet			Dimensions				
New Con	nponent / Ite	m Code	XXXXXX	Manufacturing	/ Packaging Fi	ill Site	EN-UCB Fachim SA_Chemin de Croix Blanche_Bulle_Switze
Old Com	ponent / Iten	Code	CIA79546F	Printer Name	9061245		
Pharma	Code / Laetus	Code	FPO	Print Method	Litho		
Product	/Eurocode/N	Aat. ID O	ode 8714206 8714406 73806	Substrate	white		
ArtWorks ID: EMEA 2023 00092239 000			PC spec	PC-006331			

ARTWORK SPECIFICS PANTONE PANTONE 287 C PANTONE COLUMNERT COL

DISCLAIMER - It is the client's responsibility to analyze and review immediately upon receipt of digital files, films, plates or proofs prior to any recording of matrices and/or printing. In the event of a work has been found to be incorrect after the inspection described above, contact SCHAWK to make sure that corrections are made before using through substances, have the file of the product. Schawk before the damage and the view of a model, and the substances including but not initiated to model the substance and the substances and the s

SGK is a Matthews International Corporation

FONTS USED	
HelveticaNeue MetaPlusBold-Roman	SMALLEST TEXT SIZE
MyriadPro-Regular	5.50 pt
Optima Optima-Bold	AVE. TEXT SIZE
Optima-BoldItalic ZapfDingbatsITC	7.5 pt
ZapiDingbatsire	LARGEST TEXT SIZE
	14.00 pt

If you experience any of the following, stop taking this medicine and talk to your doctor:

- Confusion
- Abnormal liver function tests
- Rapid heartbeat

Other effects which may occur include: Common: affects 1 to 10 users in 100

Headache or dizziness

- Drowsiness or tiredness
- Restlessness
- Dry mouth, sore throat
- Upset stomach, abdominal pain, diarrhoea,
- nausea

Runny nose

Uncommon: affects 1 to 10 users in 1,000

- Agitation
- Unusual weakness
- Feeling of general discomfort or uneasiness
 Pins & needles

Rarely: affects 1 to 10 users in 10,000

Aggression, depression

Weight gain

Movement disorders including tremor,

muscle spasm and ticsSleeplessness

Very rarely: affects less than 1 user in 10,000

- Taste disorder
- Blurred vision, eye movement disorder
- Difficult, painful or involuntary urination
- Other effects which may occur but it is unknown

how often:

- Vertigo, increased appetite
- Amnesia, memory impairment
- Inability to pass urine
- Eye pain
- Hepatitis
- Joint pain.
- Pruritus (intense itching) when you stop

giving this medicine.

In very rare cases people have thought about committing suicide and if you feel this way then stop taking the tablets and see your doctor. Cases of impotence in adults have also been reported.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store.

By reporting side effects you can help provide more information on the safety of this medicine.

5 Storing this medicine

Keep this medicine out of the sight and reach of children.

This medicine does not require any special storage conditions. Do not use this medicine after the expiry date stated on the carton and blister. The expiry date refers to the last day of that month. Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6 Further Information

What's in this medicine?

The active ingredient in Benadryl Allergy One a Day 10 mg Tablets is: 10 mg Cetirizine dihydrochloride in each tablet. Other ingredients are: Microcrystalline cellulose, lactose, colloidal anhydrous silica and magnesium

0£1

stearate. **The film coating contains** hydroxypropylmethylcellulose (E464), titanium dioxide (E171) and macrogol 400.

What the medicine looks like and contents of the pack

Benadryl Allergy One a Day 10 mg Tablets are white oblong film-coated tablets, with a break-line and Y-Y logo available in packs of 7, 14 and 30. Not all pack sizes may be marketed.

Product licence holder and manufacturer: McNeil Products Limited, 50 -100 Holmers Farm Way, High Wycombe, Buckinghamshire, HP12 4EG, UK.

This leaflet was revised in May 2023. Benadryl is a registered trade mark.



© McNeil Products Limited 2023

19-0127 XXXXXX