

Package Leaflet: Information for the user
Strepsils Menthol 1.2mg/0.6mg Lozenges
2,4-Dichlorobenzyl alcohol
Amylmetacresol

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.
 This medicine is available without prescription. However, you still need to take Strepsils Menthol 1.2mg/0.6mg Lozenges carefully to get the best results from it. Strepsils Menthol 1.2mg/0.6mg Lozenges will be referred to as this medicine throughout this leaflet.

- * Keep this leaflet. You may need to read it again.
- * Ask your pharmacist if you need more information or advice.
- * If you get any side effects, talk to your pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.
- * You must talk to a doctor if you do not feel better or if you feel worse after 3 days.

What is in this leaflet:
 1. What this medicine is and what it is used for
 2. What you need to know before you take this medicine
 3. How to take this medicine
 4. Possible side effects
 5. How to store this medicine
 6. Contents of the pack and other information

1. What this medicine is and what it is used for
 This medicine contains two antiseptic ingredients for use in throat medications. These are used for the symptomatic relief of mouth and throat infections. These ingredients are both mild antiseptics which kill the bacteria associated with mouth and throat infections. The action of sucking the lozenge allows the active ingredients to work in the area of the discomfort and also helps lubricate and soothe the painful area. This helps relieve the soreness and discomfort of mouth and throat infections.

2. What you need to know before you take this medicine
 This medicine is recommended for adults, children (over 12 years old) and the elderly. Children under 12 years old should not use this medicine.

Do not take this medicine if you:

- * are allergic (hypersensitive) to 2,4-dichlorobenzyl alcohol, amylmetacresol, or any of the other ingredients shown in Section 6.
- * are under 12 years old.

Warnings and precautions
Tell your doctor or pharmacist if you:

- * are taking or have recently taken any other medicines, including medicines obtained without a prescription.
- * suffer or are suffering from any other throat problems

Pregnancy and breast-feeding
Check with your pharmacist or your doctor before taking this medicine if you think you are pregnant, are pregnant or are breastfeeding.
 Ask your doctor or pharmacist for advice before taking any medicine.

Important information about some of the ingredients in this medicine

- * This medicine contains sucrose (1.38 g per lozenge) and glucose (1.10 g per lozenge).
- * This should be taken into account in patients with diabetes mellitus. If you have been told by your doctor that you have an intolerance to some sugars contact your doctor before taking this medicinal product.
- * This medicine contains only very low levels of gluten (from wheat starch). It is regarded as 'gluten-free' and is very unlikely to cause problems if you have coeliac disease. One lozenge contains no more than 22.04 micrograms of gluten. If you have wheat allergy (different from coeliac disease) you should not take this medicine.
- * This medicine contains 1.89 mg propylene glycol in each lozenge
- * This medicine contains fragrance with Benzyl Alcohol, Cinnamyl Alcohol, Citral, Citranellol, d-Limonene, Linalool (all present in Spearmint flavour), Linalool and d-limonene (present in Cool Mint Sensation flavour) and d-limonene (present in Eucalyptus oil). Benzyl Alcohol, Cinnamyl Alcohol, Citral, Citranellol, d-Limonene, Eugenol and Linalool may cause allergic reactions.



PharmacoCode reads top to bottom on the front of the medicine

PharmacoCode reads bottom to top on the back of the medicine



- * This medicine contains Sulphites – Sulphur Dioxide (E220) (present in liquid glucose) which may rarely cause severe hypersensitivity reactions and bronchospasm.

3. How to take this medicine
 Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist have told you. Check with your doctor or pharmacist if you are not sure. This medicine is for short term use only. You should take the lowest dose for the shortest time necessary to relieve your symptoms.

Adults and elderly – One lozenge to be dissolved slowly in the mouth every 2 – 3 hours up to a maximum of 12 lozenges in 24 hours
Children 12 to 16 years old – One lozenge to be dissolved slowly in the mouth every 2 – 3 hours up to a maximum of 8 lozenges in 24 hours
 Remove one lozenge from the foil blister packaging.
 Replace the foil blister back into the cardboard carton.
 Place lozenge into the mouth allowing it to dissolve slowly.
 Dissolve one lozenge slowly in the mouth every 2 – 3 hours.
 It is not recommended that children under 12 years use this medicine.
 If symptoms persist, have not improved, or have worsened after 3 days, consult a doctor or pharmacist.

If you take more of this medicine than you should
 You may experience stomach discomfort.
 Do not take any more of this medicine and consult your doctor or pharmacist.

If you forget to take this medicine
 Do not take a double dose to make up for a forgotten lozenge. Continue to use this medicine normally.
 If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects
 Like all medicines, this medicine can cause side effects, although not everybody gets them. If you experience any hypersensitivity to this medicine i.e. itching, burning sensation and/or discomfort of the mouth, difficulty in breathing, swelling of the face, neck, tongue or throat and rash (severe allergic reactions) stop taking this medicine and consult your doctor or pharmacist. If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

Reporting of side effects
 If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store this medicine
 Keep out of the sight and reach of children. Do not use this medicine after the expiry date which is stated on the carton after EXP. The expiry date refers to the last day of that month and the year. Do not store above 25°C. Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. Contents of the pack and other information
What this medicine contains:
 The active substances are: 2,4-Dichlorobenzyl alcohol 1.2mg and Amylmetacresol 0.6mg.
 The other ingredients are: Xylitol, cool mint sensation flavour (contains propylene glycol), Levomenthol, Spearmint, Eucalyptus Oil, Liquid Sucrose, Liquid Glucose (containing wheat starch (containing Gluten) and Sulphites – Sulphur Dioxide (E220)), fragrance allergens - Benzyl Alcohol, Cinnamyl Alcohol, Citral, Citranellol, d-Limonene, Eugenol and Linalool.

What this medicine looks like
 A white to pale yellow coloured lozenge with a characteristic taste of cool mint and the Strepsils brand icon pressed on both sides.

Marketing Authorisation Holder:
 Reckitt Benckiser Healthcare (UK) Limited, Slough, SL1 3UH

Manufacturer
 Reckitt Benckiser Healthcare International Limited, Nottingham, NG90 2DB
 Licence Number: PL 00063/0469
 Date of preparation: December 2020

TR2085779
 B

00000000

CUSTOMER INFO:
 Minimum Point Size = 9.00pt

Top Front

Top Back

Package Leaflet: Information for the user
Strepsils Menthol 1.2mg/0.6mg Lozenges
2,4-Dichlorobenzyl alcohol
Amylmetacresol

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.
 This medicine is available without prescription. However, you still need to take Strepsils Menthol 1.2mg/0.6mg Lozenges carefully to get the best results from it. Strepsils Menthol 1.2mg/0.6mg Lozenges will be referred to as this medicine throughout this leaflet.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get any side effects, talk to your pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.
- You must talk to a doctor if you do not feel better or if you feel worse after 3 days.

What is in this leaflet:
 1. What this medicine is and what it is used for
 2. What you need to know before you take this medicine
 3. How to take this medicine
 4. Possible side effects
 5. How to store this medicine
 6. Contents of the pack and other information

1. What this medicine is and what it is used for
 This medicine contains two antiseptic ingredients for use in throat medications. These are used for the symptomatic relief of mouth and throat infections. These ingredients are both mild antiseptics which kill the bacteria associated with mouth and throat infections. The action of sucking the lozenge allows the active ingredients to work in the area of the discomfort and also helps lubricate and soothe the painful area. This helps relieve the soreness and discomfort of mouth and throat infections.

2. What you need to know before you take this medicine
 This medicine is recommended for adults, children (over 12 years old) and the elderly. Children under 12 years old should not use this medicine.

Do not take this medicine if you:

- are allergic (hypersensitive) to 2,4-dichlorobenzyl alcohol, amylmetacresol, or any of the other ingredients shown in Section 6.
- are under 12 years old.

Warnings and precautions
Tell your doctor or pharmacist if you:

- are taking or have recently taken any other medicines, including medicines obtained without a prescription.
- suffer or are suffering from any other throat problems

Pregnancy and breast-feeding
 Check with your pharmacist or your doctor before taking this medicine if you think you are pregnant, are pregnant or are breastfeeding.
 Ask your doctor or pharmacist for advice before taking any medicine.

Important information about some of the ingredients in this medicine

- This medicine contains sucrose (1.38 g per lozenge) and glucose (1.10 g per lozenge).
- This should be taken into account in patients with diabetes mellitus. If you have been told by your doctor that you have an intolerance to some sugars contact your doctor before taking this medicinal product.
- This medicine contains only very low levels of gluten (from wheat starch). It is regarded as 'gluten-free' and is very unlikely to cause problems if you have coeliac disease. One lozenge contains no more than 22.04 micrograms of gluten. If you have wheat allergy (different from coeliac disease) you should not take this medicine.
- This medicine contains 1.89 mg propylene glycol in each lozenge
- This medicine contains fragrance with Benzyl Alcohol, Cinnamyl Alcohol, Citral, Citranellol, d-Limonene, Linalool (all present in Spearmint flavour), Linalool and d-limonene (present in Cool Mint Sensation flavour) and d-limonene (present in Eucalyptus oil). Benzyl Alcohol, Cinnamyl Alcohol, Citral, Citranellol, d-Limonene, Eugenol and Linalool may cause allergic reactions.

- This medicine contains Sulphites – Sulphur Dioxide (E220) (present in liquid glucose) which may rarely cause severe hypersensitivity reactions and bronchospasm.

3. How to take this medicine
 Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist have told you. Check with your doctor or pharmacist if you are not sure. This medicine is for short term use only. You should take the lowest dose for the shortest time necessary to relieve your symptoms.

Adults and elderly – One lozenge to be dissolved slowly in the mouth every 2 – 3 hours up to a maximum of 12 lozenges in 24 hours
Children 12 to 16 years old – One lozenge to be dissolved slowly in the mouth every 2 – 3 hours up to a maximum of 8 lozenges in 24 hours
 Remove one lozenge from the foil blister packaging.
 Replace the foil blister back into the cardboard carton.
 Place lozenge into the mouth allowing it to dissolve slowly.
 Dissolve one lozenge slowly in the mouth every 2 – 3 hours.
 It is not recommended that children under 12 years use this medicine.
 If symptoms persist, have not improved, or have worsened after 3 days, consult a doctor or pharmacist.

If you take more of this medicine than you should
 You may experience stomach discomfort.
 Do not take any more of this medicine and consult your doctor or pharmacist.

If you forget to take this medicine
 Do not take a double dose to make up for a forgotten lozenge. Continue to use this medicine normally.
 If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects
 Like all medicines, this medicine can cause side effects, although not everybody gets them. If you experience any hypersensitivity to this medicine i.e. itching, burning sensation and/or discomfort of the mouth, difficulty in breathing, swelling of the face, neck, tongue or throat and rash (severe allergic reactions) stop taking this medicine and consult your doctor or pharmacist. If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

Reporting of side effects
 If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store this medicine
 Keep out of the sight and reach of children. Do not use this medicine after the expiry date which is stated on the carton after EXP. The expiry date refers to the last day of that month and the year. Do not store above 25°C Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. Contents of the pack and other information
What this medicine contains:
 The active substances are: 2,4-Dichlorobenzyl alcohol 1.2mg and Amylmetacresol 0.6mg
 The other ingredients are: Xylitol, cool mint sensation flavour (contains propylene glycol), Levomenthol, Spearmint, Eucalyptus Oil, Liquid Sucrose, Liquid Glucose (containing wheat starch (containing Gluten) and Sulphites – Sulphur Dioxide (E220)), fragrance allergens - Benzyl Alcohol, Cinnamyl Alcohol, Citral, Citranellol, d-Limonene, Eugenol and Linalool.
What this medicine looks like
 A white to pale yellow coloured lozenge with a characteristic taste of cool mint and the Strepsils brand icon pressed on both sides.


Marketing Authorisation Holder:
 Reckitt Benckiser Healthcare (UK) Limited, Slough, SL1 3UH
Manufacturer
 Reckitt Benckiser Healthcare International Limited, Nottingham, NG90 2DB
 Licence Number: PL 00063/0469
 Date of preparation: December 2020

TR2204988
 B

Pharmacode Reads This Way ↑
 Last Bar
 First Bar
 ↓
 72.5mm from first bar to bottom of leaflet

31889903

83.6mm to centre of mark



CUSTOMER INFO:
 Minimum Point Size = 9.00pt